ELSEVIER

Toxicology 126 (1998) 239–240

TOXICOLOGY

Volume contents

Volume 126 (1998)

Metabolism and toxicity of diisopropylnaphthalene as compared to naphthalene and monoalkyl naphthalenes: a minireview
There is an inverse relation between metabolic conversion and toxicity
H. Höke, R. Zellerhoff (Germany) 1–7

Altered zinc metabolism contributes to the developmental toxicity of 2-ethylhexanoic acid, 2-ethylhexanol and valproic acid
The teratogenic effects of the chemicals can be modulated by dietary zinc intake
L.M. Bui, M.W. Taubeneck, J.F. Commisso, J.Y. Uriu-Hare, W.D. Faber, C.L. Keen (USA). 9–21

Decreased hepatobiliary secretion of inorganic mercury, its deposition and toxicity in the Eisai hyperbilirubinemic rat with no hepatic canalicular organic anion transporter
The mutant rats also have kidneys highly sensitive to Hg poisoning
N. Sugawara, Y.-R. Lai, C. Sugaware, K. Arizono (Japan). 23–31

Cutaneous tolerance to nitroxide free radicals and nitrone spin traps in the guinea pig
The tested nitroxides and nitrones show low potential for acute skin intolerance
J. Fuchs, N. Groth, T. Herrling (Germany) 33–40

Augmentation of allergic reactions by several pesticides
Test systems were histamine release from mast cells in vitro and passive cutaneous anaphylaxis in vivo
T. Sato, M. Taguchi, H. Nagase, H. Kito, M. Niikawa (Japan) 41–53

γ-Glutamylcysteinylethyl ester attenuates progression of carbon tetrachloride-induced acute liver injury in mice
Progression may be attenuated by maintenance of hepatic glutathione levels
K. Nishida, Y. Ohta, I. Ishiguro (Japan). 55–63

Effect of the peroxisome proliferator ciprofibrate on hepatic cyclooxygenase and phospholipase A_2 in rats
Ciprofibrate increases the expression of cyclooxygenase 2 protein
L.K. Leung, H.P. Glauert (USA) 65–73

The guinea-pig skin sensitization test revisited: an evaluation formula to predict possible sensitization levels for eight chemicals used in household products
A new formula is proposed to assess the sensitizing potential of chemicals
J. Momma, S. Kitajima, T. Inoue (Japan) 75–82

Lipid peroxidation of rat myocardial tissue following daunomycin administration
Increased lipid peroxidation is discussed as a toxic mechanism
G. Papageorgiou, S. Iliadis, N. Botsoglou, C. Dioudis, A. Goulas, D. Fletouris, A. Dimitriadou-Vafiadou (Greece). 83–91

DNA oxidation by potassium bromate; a direct mechanism or linked to lipid peroxidation?
The answer to the question seems to be complex, according to the doses applied
J.K. Chipman, J.E. Davies, J.L. Parsons, J. Nair, G. O'Neill, J.K. Fawell (UK, Germany) 93–102

Evaluation of the direct genotoxic potential of cadmium in four different rodent cell lines
Increased DNA damage was only observed at Cd concentrations that arrested cell growth
R.R. Misra, G.T. Smith, M.P. Waalkes (USA) 103–114

Kinetics of aluminum disposition after ingestion of low to moderate pharmacological doses of aluminum.
Doses of Al comparable to uptake by humans in buffered aspirin or antacids were studied
J.E. Sutherland, J.L. Greger (USA). 115–125

The enzyme inducers 3-methylcholanthrene and phenobarbital affect the activities of glucocorticoid hormone-regulated enzymes in rat liver and kidney
M. Boll, L.W. Weber, M. Font, A. Stampfl (Germany, USA, Spain) . . . . . .
Induction of various cytosolic enzymes in a peculiar fashion with two maxima in the time curve 127–136
Further evaluation of the incorporation of an immunotoxicological functional assay for assessing humoral immunity for hazard identification purposes in rats in a standard toxicology study
Injection of sheep red blood cells for assessing humoral immunity did not alter standard toxicological endpoints
G.S. Ladics, C. Smith, G.S. Elliott, T.W. Slone, S.E. Loveless (USA). 137–152

Announcement 153

Glutathione *S*-transferase Mu (GST Mu) deficiency and DNA adducts in lymphocytes of smokers
Individuals deficient in GST MU may be more susceptible to smoke-induced DNA damage
M. Soni, M. Madurantakan, K. Krishnaswamy (India) 155–162
Glutathione *S*-transferase Mu (GST Mu) deficiency and DNA adducts in lymphocytes of smokers
Sulfhydryl groups appear to play an essential role in coproporphyrinogen
M. Soni, M. Madurantakan, K. Krishnaswamy (India) 163–171
Lack of carcinogenicity of cadmium chloride in female Syrian hamsters
Three different single doses of cadmium injected s.c followed by life time observation
M.P. Waalkes, S. Rehm (USA) 173–178
Marked inhibition of hepatic cytochrome P450 activity in cholesterol-induced atherosclerosis in rabbits
One percent cholesterol feeding leads to diversified effects on durg metabolising activities and the reapective apoproteins
A. Irizar, C. Ioannides (UK). 179–193
Concurrent administration of D-penicillamine and zinc has no advantages over the use of either single agent on copper excretion in the rat
Combination of both therapeutic principles for Wilson's disease offers no advantage over single treatment
J.L. Domingo, M. Gómez, M.M. Jones (Spain, USA) 195–201
Alterations of thiol metabolism in human cell lines induced by low amounts of copper, mercury or cadmium ions
Copper affects thiols and releases homocysteine, the others increase glutathione synthesis
B. Hultberg, A. Andersson, A. Isaksson (Sweden) 203–212
Protective effects of aged garlic extract against bromobenzene toxicity to precision cut rat liver slices
Addition of one to five percent extract for several hours improves various parameters of cytotoxicity
B.H. Wang, K.A. Zuzel, K. Rahman, D. Billington (UK). 213–222
Renal tolerance of digoxin-specific Fab fragments in the rat
No renal nephrotoxicity was observed when administering digoxinspecific Fab fragments to rats
E. Advenier, S. Mouajjah, M. Bazin-Redureau, J.-M. Launay, J.-M. Scherrmann (France) 223–229
Vth Conference of the International Society for Trace Element Research in Humans (ISTERH): New aspects of trace element research. 231

Author index. 233
Subject index. 235–238
Volume contents. 239–240